arXiv:1212.1526v1 [math.CV] 7 Dec 2012

Integral-type operators from Hardy space to Bloch space on the upper half-plane

Ning Xu *

Department of Mathematics and Science, Huai Hai Institute of Technology,

Jiangsu, Lianyungang 222005, P. R. China

Abstract. The boundedness of compactness of integral-type operators from Hardy space to Bloch space on the upper half-plane $\Pi_+ = \{z \in \mathbb{C} : Imz > 0\}$ are characterized.

Keywords: Integral-type operator Upper half-plane Boundedness Compactness

1. Introduction

Let Π_+ be the upper half-plane and $H(\Pi_+)$ be the space of all holomorphic functions on the Π_+. The Hardy spaces $H^2(\Pi_+)$ consists of all $f \in H(\Pi_+)$ such that

$$\|f\|^2_{H^2(\Pi_+)} = \sup_{y>0} \int_{-\infty}^{\infty} |f(x+iy)|^2 dx < \infty$$

where $y = Imz$.

The Bloch space $B_\infty(\Pi_+)$ consists of all of $f \in H(\Pi_+)$ such that

$$B(f) = \sup_{z \in \Pi_+} Imz|f'(z)| < \infty.$$

A natural norm on the Bloch space can be introduced as follows

$$\|f\|^2_{B_\infty(\Pi_+)} = |f(i)| + B(f).$$

With this norm the $B_\infty(\Pi_+)$ becomes a Banach space.

For $f \in H(\Pi_+)$,fix $x_0 \in \Pi_+$ the integral-type operators J_g and I_g are defined by

$$J_g f(z) = \int_{z_0}^{z} f(\zeta) g'(\zeta) d\zeta, z \in \Pi_+ \qquad (1)$$

---

[*] Corresponding author, e-mail: gx899200@126.com.

$$I_g f(z) = \int_{z_0}^{z} f'(\zeta)g(\zeta)d\zeta, z \in \Pi_+ \qquad (2)$$

where $g \in H(\Pi_+)$.

The boundedness and compactness of J_g and I_g have been introduced on the unit dick in[1,8,10,11,20].The operators J_g and I_g, as well as their n-dimensional generalizations,acting on various spaces of analytic functions, have been recently studied,for example, in[2-7,9,19].

The importance of the operators J_g and I_g comes from that

$$J_g f + I_g f = M_g f - f(z_0)g(z_0)$$

where the multiplication operator M_g is defined by $(M_g f)(z) = g(z)f(z)$.

Let X and Y be topological vector spaces whose topologies are given by translation- invariant metrics d_X and d_Y,respectively.It is said that a linear operator $T : X \to Y$ is metrically bounded if there exists a positive constant K such that

$$d_Y(Tf, 0) \leq d_X(f, 0)$$

for all $f \in X$. When X and Y are Banach spaces, the metrically boundedness coincides with the boundedness of operators between Banach spaces. If we say that an operator is bounded we will regard that it is metrically bounded. Operator $T : X \to Y$ is said to be metrically compact if it takes every metric ball in X into a relatively compact set in Y.

While there is a vast literature on composition and weighted composition operators between spaces of holomorphic functions on the unit disk,there are few papers on there operators on spaces of functions holomorphic on the upper half-plane(see,12-18). In this paper,we consider the metrically boundedness and compactness of (1),(2) acting from $H^2(\Pi_+)$ to $B_\infty(\Pi_+)$ on the upper half-plane.

Throughout this paper, constants are denoted by C,they are positive and may differ from one occurrence to the other. The notion $A \asymp B$ means that there is a positive constant C such that $C^{-1}B \leq A \leq CB$.

2. Auxiliary results

Lemma 1. Let $L = J_g(I_g)$ and $g \in H^\infty(\Pi_+)$. Then$L : H^2(\Pi_+)$ to $B_\infty(\Pi_+)$ is metrically compact if and only if for any bounded sequence $\{f_n\}_{n\in\mathbb{C}}$ in $H^2(\Pi_+)$ converging to zero on compact of Π_+, we have $\lim_{n\to\infty} \|Lf_n\|_{B_\infty(\Pi+)} = 0$.

The proof is standard which can be found in [14] and omitted here.

Lemma 2[14]**.**Let $0 < p < \infty$ and $0 < a < b$. If $f \in H^2(\Pi_+)$, then

$$\lim_{z\to\infty z\in\Gamma_{a,b}} f(z) = 0,$$

where $\Gamma_{a,b} = \{z \in \Pi_+ : a \leq Imz \leq b\}$.

Lemma 3[17]**.** Let $f \in H^2(\Pi_+)$ then

$$|f^{(n)}(z)| \leq C\frac{\|f\|_{H^2(\Pi_+)}}{(Imz)^{n+\frac{1}{2}}}, n = 0, 1, 2 \qquad (3)$$

3.Main results

Here we formulate and prove the main results of this paper.

Theorem 1. Let g is holomorphic function of Π_+ and $g \in H^\infty(\Pi_+)$. Then

(a) $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded if and only if

$$M_1 = \sup_{z\in\Pi_+} (Imz)^{\frac{1}{2}}|g'(z)| < \infty. \qquad (4)$$

(b) $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded if and only if

$$M_2 = \sup_{z\in\Pi_+} \frac{|g(z)|}{(Imz)^{\frac{1}{2}}} < \infty. \qquad (5)$$

Proof.(a)Assume that condition in(4) hold. Then

$$\begin{aligned} \|J_g f\|_{B_\infty(\Pi_+)} &= |J_g f(i)| + \sup_{z\in\Pi_+} Imz|f(z)g'(z)| \\ &= |\int_{z_0}^{i} f(\zeta)g'(\zeta)d\zeta| + \sup_{z\in\Pi_+} Imz|f(z)g'(z)| \\ &\leq \max_{|z_0|\leq|\zeta|\leq 1, \zeta\in\Pi_+} |f(\zeta)||g(i) - g(z_0)| \\ &+ \sup_{z\in\Pi_+} (Imz)^{\frac{1}{2}}|g'(z)|\|f\|_{H^2(\Pi_+)}. \end{aligned}$$

From $f \in H^2(\Pi_+)$ and $g \in H^\infty(\Pi_+)$, it follows that $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded.

Now assume that $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded. Choose

$$f_w(z) = \frac{(Imw)^{\frac{3}{2}}}{\Pi^{\frac{1}{2}}(z - \bar{w})^2} \qquad (6)$$

It is clear that $f_w \in H^2(\Pi_+)$ and that $\|f_w\|_{H^2(\Pi_+)} = 1$. Choose $w = z$, we obtain

$$C \geq \|J_g f_w\|_{B_\infty(\Pi_+)} \geq \sup_{w\in\Pi_+} Imw|f_w(w)g'(w)| = \sup_{w\in\Pi_+} \frac{(Imz)^{\frac{1}{2}}|g'(z)|}{4\Pi^{\frac{1}{2}}}$$

Therefore we obtain (4).

(b) Assume that condition in (5) hold. Then

$$\begin{aligned}\|I_gf\|_{B_\infty(\Pi_+)} &= |I_gf(i)| + \sup_{z\in\Pi_+} Imz|f'(z)g(z)| \\ &= |\int_{z_0}^{i} f'(\zeta)g(\zeta)d\zeta| + \sup_{z\in\Pi_+} Imz|f'(z)g(z)| \\ &\leq |f(i)-f(z_0)| \max_{|z_0|\leq|\zeta|\leq 1, \zeta\in\Pi_+} |g(\zeta)| \\ &+ \sup_{z\in\Pi_+} (Imz)^{\frac{1}{2}}|g'(z)|\|f\|_{H^2(\Pi_+)}.\end{aligned}$$

From $f \in H^2(\Pi_+)$ and $g \in H^\infty(\Pi_+)$, it follows that $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded.

Now assume that $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded. Choose(6) and let $w = z$, we obtain

$$C \geq \|I_gf_w\|_{B_\infty(\Pi_+)} \geq \sup_{w\in\Pi_+} Imw|f'_w(w)g(w)| = \sup_{w\in\Pi_+} \frac{|g(w)|}{4\pi^{\frac{1}{2}}(Imw)^{\frac{1}{2}}}.$$

Therefore we obtain (5).

Theorem 2. Let g is holomorphic function of Π_+ and $g \in H^\infty(\Pi_+)$.

(a) $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact, then

$$\lim_{r\to 0} \sup_{y<r} (Imz)^{\frac{1}{2}}|g'(z)| = 0. \qquad (7)$$

(b) $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact, then

$$\lim_{r\to 0} \sup_{y<r} \frac{|g(z)|}{(Imz)^{\frac{1}{2}}} = 0. \qquad (8)$$

where it is understood that if $\{z : Imz < r\}$ is empty for some $r > 0$, the supremum is equal to zero.

Proof.(a) Suppose that $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact and (7) does not hold. Then there exists a positive number δ and a sequence $\{z_n\}_{n\in\mathbb{N}}$ in Π_+ such that $Imz_n \to 0$ and

$$(Imz_n)^{\frac{1}{2}}|g'(z_n)| > \delta, \quad n \in \mathbb{N}.$$

Choose

$$f_n(z) = \frac{(Imw_n)^{\frac{3}{2}}}{\Pi^{\frac{1}{2}}(z-\bar{w_n})^2} \qquad (9)$$

and let $z_n = w_n$,then f_n is norm bounded in $H^2(\Pi_+)$ and $f_n \to 0$ uniformly on compacts of Π_+ as $Imz_n \to 0$. By Lemma 2 it follows that a subsequence of $\{J_g f_n\}$ tends to 0 in $B_\infty(\Pi_+)$. On the other hand,

$$\|J_g f\|_{B_\infty(\Pi_+)} \geq Imw_n|f_n(w_n)g'(w_n)| = \frac{1}{\sqrt{\pi}}(Imw_n)^{\frac{1}{2}}|g'(w_n)| = \frac{\delta}{\sqrt{\pi}},$$

which is a contradiction.

(b) Suppose that $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact and (8) does not hold. Then there exists a positive number δ and a sequence $\{z_n\}_{n\in\mathbb{N}}$ in Π_+ such that $Imz_n \to 0$ and

$$\frac{|g(z_n)|}{(Imz_n)^{\frac{1}{2}}} > \delta$$

Choose (9) and let $z_n = w_n$, we obtain

$$\|I_g f\|_{B_\infty(\Pi_+)} \geq Imw_n|f_n'(w_n)g(w_n)| = \frac{1}{4\sqrt{\pi}}\frac{|g(w_n)|}{(Imw_n)^{\frac{1}{2}}} = \frac{\delta}{4\sqrt{\pi}},$$

which is a contradiction.

Theorem 3. Let$g$ is holomorphic function of Π_+ ,$g \in H^\infty(\Pi_+)$ and $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is bounded. Suppose that $g \in B_\infty(\Pi_+)$. Then

(a) $J_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact if

$$\lim_{r\to 0}\sup_{y<r}(Imz)^{\frac{1}{2}}|g'(z)| = 0. \tag{10}$$

(b) $I_g : H^2(\Pi_+) \to B_\infty(\Pi_+)$ is compact if

$$\lim_{r\to 0}\sup_{y<r}\frac{|g(z)|}{(Imz)^{\frac{1}{2}}} = 0. \tag{11}$$

Proof.(a) Assume that (10) holds. Then for every $\epsilon > 0$, there exists an $M_1 > 0$ such that

$$\sup_{y<r}(Imz)^{\frac{1}{2}}|g'(z)| < \epsilon, \quad whenever Imz < M_1. \tag{12}$$

Assume $\{f_n\}_{n\in\mathbb{N}}$ is a sequence in $H^2(\Pi_+)$ such that $\sup_{n\in\mathbb{N}}\|f_n\|_{H^2(\Pi_+)} \leq M$ and $f_n \to 0$ uniformly on compacts of Π_+ as $n \to \infty$. Thus for $z \in \Pi_+$,such that $Imz < M_1$ and each $n \in \mathbb{N}$, we have

$$Imz|f_n(z)g'(z)| \leq C(Imz)^{\frac{1}{2}}|g'(z)|\|f_n\|_{H^2(\Pi_+)} \leq \epsilon CM. \tag{13}$$

By(3), we have

$$|f_n(z)| \leq C\frac{\|f_n\|_{H^2(\Pi_+)}}{(Imz)^{\frac{1}{2}}} \leq C\frac{M}{(Imz)^{\frac{1}{2}}}.$$

Thus there is an $M_2 > M_1$ such that

$$|f_n(z)| < \epsilon, \quad Whenever Imz > M_2.$$

Hence for $z \in \Pi_+$ such that $Imz > M_2$,and each $n \in \mathbb{N}$, we have

$$(Imz)|f_n(z)g'(z)| < \epsilon\|g\|_{B^{\infty}(\Pi_+)}. \qquad (14)$$

If $M_1 \leq Imz \leq M_2$, then by Lemma 2,there exists an $M_3 > 0$ such that

$$|f_n(z)| < \epsilon, \quad whenever |Rz| > M_3.$$

Therefore, for each $n \in \mathbb{N}$, when $|Rz| > M_3$, we have

$$Imz|f_n(z)g'(z)| < \epsilon\|g\|_{B^{\infty}(\Pi_+)}. \qquad (15)$$

If $M_1 \leq Imz \leq M_2$ and $|Rz| \leq M_3$, then there exists some $n_0 \in \mathbb{N}$ such that $|f_n(z)| < \epsilon$ for all $n \geq n_0$ and so

$$Imz|f_n(z)g'(z)| < \epsilon\|g\|_{B^{\infty}(\Pi_+)}. \qquad (16)$$

Finally, we also have

$$|J_g f_n(i)| = |\int_{z_0}^{i} f_n(\zeta)g'(\zeta)d\zeta| \leq \max_{|z_0|\leq|\zeta|\leq 1, \zeta\in\Pi_+} |f_n(\zeta)||g(i) - g(z_0)| \quad (17)$$

as $n \to \infty$.

Combining (12)-(17), we have$\|J_g f_n\|_{B_{\infty}(\Pi_+)} < \epsilon C$, for $n \geq n_0$ and some $C > 0$. Thus by Lemma 1,$J_g : H^2(\Pi_+) \to B_{\infty}(\Pi_+)$ is compact.

REFERENCES

1. A.Aleman, A.G.Siskakis,An integral operator on H^p,Complex Variables Theory and Application,28(2)(1995)149-158.

2. A.Aleman,J.A.Cima,An integral operator on H^p and Hardy's inequality, J.Anal.Math.85(2001)157-176.

3. D.C. Chang,S.Li,S.Stević,On some integral operators on the unit polydisk and the unit ball, Taiwanese J. Math.11(5)(2007)1251-1286.

4. D.C. Chang, S.Stević, Estimates of an integral operator on function spaces, Taiwanese J. Math. 7(3)(2003)423-432.

5. D.C. Chang, S.Stević,The generalized Cesàro operator on the unit polydisk, Taiwanese J. Math.7(2)(2003)293-308.
6. Z.Hu, Extended Cesàro operators on the Bloch spaces in the unit ball of $\mathbb{C}^n$, Acta. Math.Sci.Ser.B Engl. Ed.23(4)(2003)561-566.
7. Z.Hu, Extended Cesàro operators on mixed norm spaces, Proc. Amer. Math. Soc. 131(7)(2003)2171-2179.
8. S.Li,S.Stević, Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to the Zygmund space,J.Math.Anal.Appl.345 (2008)40-52.
9. S. Li, S.Stević,Cesàro-type operators on some spaces of analytic functions on the unit ball,Appl.Math.Comput. 208(2009)378-388.
10. S. Li, S.Stević,Products of integral-type operators and composition operators between Bloch-type spaces, J.Math.Anal.Appl.349(2009)596-610.
11. S. Li, S.Stević,Volterra-type Operators on Zygmund spaces,J.Inequal.Appl.(1) (2005)81-88.
12. V.Matache,Compact composition operators on H^p of a half-plane, Proc. Amer. Math. Soc. 127(1999)1483-1491.
13. S.D.Sharma,A.K.Sharma,S.Ahmed,Composition operators between Hardy and Bloch-type spaces of the upper half-plane, Bull.Korean Math.Soc.44(2007)475-482.
14. S.Stević,Ajay K.Sharma,Weighted composition operators between Hardy and growth spaces on the upper half-plane, Appl.Math.Comput,217(2011)4928-4934.
15. S.Stević,Composition operators from the weighted Bergman space to the nth weighted-type space on the upper half-plane, Appl.Math.Comput,217 (2010)3379-3384.
16. S.Stević,Composition operators from the Hardy space to the Zygmund-type space on the upper half-plane, Abstr. Appl. Anal.,2009(2009).Article ID161528,8 pages.

17. S.Stević,Composition operators from the Hardy space to the nth weighted-type space on the unit disk and the half-plane, Appl.Math.Comput,215(2010)3950-3955.

18. S.Stević,Composition operators from the Hardy space to the Zygmund-type space on the upper half-plane and the unit disk, J.Comput.Anal.Appl.12(2)(2010)305-312.

19. N.Xu, Extended Cesàro operators on μ-Bloch spaces in $\mathbb{C}^n$, J. Math. Research and Exposition, 29(5)(2009)913-922.

20. W.Yang,On an integral-type operator between Bloch-type spaces,Appl.Math. Comput,215(3)(2009)954-960.